

19



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

11 Publication number:

0 368 486  
A2

12

## EUROPEAN PATENT APPLICATION

21 Application number: 89310620.3

22 Date of filing: 17.10.89

51 Int. Cl.<sup>5</sup>: C07K 5/08, C07K 7/06,  
C07K 7/10, C12P 21/00,  
C12N 15/00, A61K 37/02,  
C07K 13/00

30 Priority: 10.11.88 US 269583

43 Date of publication of application:  
16.05.90 Bulletin 90/20

64 Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: MERCK & CO. INC.  
126, East Lincoln Avenue P.O. Box 2000  
Rahway New Jersey 07065-0900(US)

Applicant: THE TRUSTEES OF THE  
UNIVERSITY OF PENNSYLVANIA  
c/o Office of Research Administration 133  
South 36th Street Suite 300  
Philadelphia Pennsylvania 19104-3246(US)

72 Inventor: Shattil, Sanford J.  
510 Merwyn Road  
Narberth Pennsylvania 19072(US)  
Inventor: Taub, Rebecca A.  
403 Great Spring Road  
Rosemont Pennsylvania 19010(US)  
Inventor: Friedman, Paul A.  
403 Great Spring Road  
Rosemont Pennsylvania 19010(US)

74 Representative: Stebbing, Peter John Hunter  
Stebbing & Co. 45 Lancaster Mews  
Lancaster Gate, London W2 3QQ(GB)

54 Novel inhibitors of platelet binding.

57 Peptides comprising the tripeptide sequence of arginine-tyrosine-aspartic acid, linked from amino- to carboxy-terminus, are inhibitors of fibrinogen binding, platelet aggregation, glycoprotein IIb-IIIa binding and related aberrant and normal physiological conditions.

EP 0 368 486 A2

## Novel Inhibitors of Platelet Binding

### BACKGROUND OF THE INVENTION

Many adhesive proteins, including fibrinogen, fibronectin, von Willebrand factor, and vitronectin, contain an amino acid sequence that functions as a recognition site for adhesion receptors. The amino acid sequences within these proteins are recognized by structurally-related receptors on a variety of mammalian cells that have been termed integrins or cytoadhesions. Hynes, *Cell*, 48, 549-554 (1987); Ruoslahti et al., *Science*, 238, 491-497 (1987).


One well-characterized member of the integrin family is the blood platelet membrane glycoprotein IIb-IIIa complex (GP IIb-IIIa). Phillips et al., *Blood*, 71, 831-843 (1988). Upon platelet activation, GP IIb-IIIa becomes competent to bind fibrinogen, a process required for platelet aggregation. Bennett et al., *J. Clin. Invest.*, 64, 1393-1400 (1979); Marguerie et al., *J. Biol. Chem.*, 254, 5357-5363 (1979). Based on studies with synthetic peptides, the Arg-Gly-Asp-Ser sequence at positions 572-575 of the fibrinogen A $\alpha$  chain appears to play a major role in the interaction of this protein with GP IIb-IIIa, although other regions in fibrinogen may be involved as well. Gartner et al., *J. Biol. Chem.*, 260, 11891-11894 (1985); Ginsberg et al., *J. Biol. Chem.*, 260, 3931-3936 (1985); Haverstick et al., *Blood*, 66, 946-952 (1985); Pytela et al., *Science*, 231, 1159-1162 (1986); Kloczewiak et al., *Thrombosis Res.*, 29, 249-255 (1983); Kloczewiak et al., *Biochemistry*, 23, 1767-1774 (1984). However, the precise details of the molecular interaction between fibrinogen and GP IIb-IIIa are, at present, unknown. The tripeptide sequence Arg-Gly-Asp has also been identified in the adhesive proteins fibronectin, von Willebrand factor and vitronectin. Hynes, *Cell*, 48, 549-554 (1987); Ruoslahti et al., *Science*, 238, 491-497 (1987). A similar region of fibronectin containing the sequence Arg-Glu-Asp-Val has been implicated in the binding of this protein to melanoma cells. Humphries et al., *J. Cell Biol.*, 103, 2637-2647 (1986). Moreover, von Willebrand factor, which can bind to GP IIb-IIIa on activated platelets, contains, in its long (significantly greater than 50 residue) amino acid sequence, an Arg-Tyr-Asp-Ala, in addition to an Arg-Gly-Asp-Ser, sequence. Titani et al., *Biochemistry*, 25, 3171-3184 (1986). Furthermore, the sequence Arg-Phe-Asp-Ser and Arg-Tyr-Asp-Ser are found within the long chain peptides that comprise the major histocompatibility antigens, peptides which have a length greatly in excess of 50 amino acid residues. Auffray et al., *Human Immunol.*, 15, 381-390 (1986).

Numerous applications have been found for peptides containing the amino acid sequence Arg-Gly-Asp. The binding of fibrinogen to blood platelets, for example, has been found to be inhibited by Arg-Gly-Asp-containing peptides. In addition, PAC1, an IgM-x murine monoclonal antibody that, like fibrinogen, appears to bind to GP IIb-IIIa on activated platelets, is similarly inhibited by such peptides. The inhibition of fibrinogen and PAC1 occurs at an apparent  $K_i$  of about 10-20  $\mu$ M when the tetrapeptide Arg-Gly-Asp-Ser is employed. Shattil et al., *J. Biol. Chem.*, 260, 11107-11114 (1985); Bennett et al., *J. Biol. Chem.*, (1988); Shattil et al., *Blood*, 68, 1224-1231 (1986). Moreover, U.S. Patent No. 4,683,291 discloses the usefulness of peptides containing the Arg-Gly-Asp sequence in inhibiting cellular adhesion in general, and platelet to platelet binding specifically. Such peptides are also reported as useful in retarding the formation of blood clots.

Additional and/or better peptides which meet the important ends of inhibiting fibrinogen to platelet binding, platelet to platelet binding and/or blood clot formation are needed. The present invention is directed to these and other important needs.

### SUMMARY OF THE INVENTION

The present invention provides compounds of the general Formula I



and physiologically acceptable salts thereof, wherein:


$X^1$  and  $X^2$ , independently, are 1 to about 50 amino acid residues, provided that the total number of amino acid residues in  $X^1$  and  $X^2$ , taken together, is less than or equal to about 50;

$m$  and  $n$ , independently, are 0 or 1;

$R^1$  is a protected or unprotected terminal amino group; and

$R^2$  is a protected or unprotected terminal carboxy group.

Also included in the subject invention are compounds of the general Formula II



and physiologically acceptable salts thereof, wherein:

$X^1$  and  $X^2$ , independently, are 1 or more amino acid residues;

$Y$  is 1 or more amino acid residues, provided that  $Y$  does not include a Ser or Ala residue;

$m$ ,  $n$  and  $o$ , independently, are 0 or 1, provided that when  $m$  is 1,  $o$  must be 1;

R<sup>1</sup> is a protected or unprotected terminal amino group; and

R<sup>2</sup> is a protected or unprotected terminal carboxy group; and wherein said compound inhibits fibrinogen binding, platelet aggregation, glycoprotein IIb-IIIa binding, thrombosis and/or cancer metastasis.

The invention further provides compositions comprising one or more of the foregoing compounds, and methods of using such compounds or compositions in inhibiting fibrinogen binding, platelet aggregation, and/or glycoprotein IIb-IIIa binding. The subject compounds or compositions are also effective in the treatment of certain physiological conditions, such as thrombosis and/or cancer metastasis, and the present invention is further directed to these uses.

Also included within the ambit of the present invention are monoclonal and polyclonal antibodies directed to the subject compounds and nucleic acid fragments comprising nucleotide sequences which encode the subject compounds.

Finally, the present invention comprehends a peptide comprising an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\mu$  heavy chain of monoclonal antibody PAC1, as well as antibodies incorporating that sequence. The present invention further includes a peptide comprising an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\mu$  heavy chain of monoclonal antibody PAC1, and an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\mu$  light chain of monoclonal antibody PAC1, as well as antibodies incorporating that sequence. Most specifically, the invention is directed to the monoclonal antibody PAC1.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of the variable region of the  $\kappa$  light chain of a PAC1 hybridoma cell, with nucleotide numbering shown above. The site of the V to J region junction is indicated. The predicted amino acid sequence, along with the amino acid numbers appear on the next two lines below. The amino acids are numbered according to convention. Kabat E.A., Wu, T.T., Reid-Miller, M., Petty, H.M., Gottesman, K.S., Sequences of Proteins of Immunologic Interest (US DHHS, PHS, NIH 1987). The complimentary determining regions (CDRs) are shown in boxes.

Figure 2 shows the nucleotide sequence of the variable region of the  $\mu$  heavy chain of a PAC1 hybridoma cell with nucleotide numbering shown

above. The site of the V to J region junction is indicated. The predicted amino acid sequence, along with the amino acid numbers appear on the next two lines below. The amino acids are numbered according to convention. Kabat E.A., Wu, T.T., Reid-Miller, M., Perry, H.M., Gottesman, K.S., Sequences of Proteins of Immunologic Interest (US DHHS, PHS, NIH 1987). The complimentary determining regions (CDRs) are shown in boxes. CDR3 is comprised of the amino acid sequence Arg-Ser-Pro-Ser-Tyr-Tyr-Arg-Tyr-Asp-Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr.

Figure 3 shows the effect of a 21 amino acid peptide derived from the variable region of the PAC1  $\mu$  heavy chain (hereinafter referred to as "the Arg-Tyr-Asp-21mer peptide") on the binding of PAC1 and fibrinogen to activated human blood platelets. The Arg-Tyr-Asp-21mer has the sequence Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr-Arg-Tyr-Asp-Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr. Gel-filtered platelets were incubated for 15 min with ADP (10  $\mu$ M), and epinephrine (10  $\mu$ M), in the presence of various peptides and either FITC-PAC1 (Figure 3A) or FITC-9F9 (Figure 3B) and fibrinogen (50  $\mu$ g/ml). The specific peptides tested were Arg-Tyr-Asp-21mer (filled circles), Arg-Gly-Asp-Ser (unfilled circles) and an unrelated-28mer (control; unfilled squares). Binding was determined by flow cytometry. Antibody binding is expressed as the percent of binding observed in the absence of added peptide. Error bars represent the mean  $\pm$  SEM of 5 separate experiments. In Figure 2A, the data points without error bars represent the mean of two experiments.

Figure 4 shows the effect of the Arg-Tyr-Asp-21mer peptide on the binding of various monoclonal antibodies to platelets. Gel-filtered platelets were incubated in the presence of Arg-Tyr-Asp-21mer and various FITC-labelled anti-platelet monoclonal antibodies. Monoclonal antibody binding was determined by flow cytometry. In the case of antibody AP1 (specific for platelet glycoprotein Ib) (unfilled circles), the platelets were not activated. In the case of antibodies PAC1 (specific for activated glycoproteins IIb-IIIa) (filled circles), B1B5 (specific for glycoprotein IIb) (filled squares) and S12 (specific for GMP-140) (unfilled triangles), the platelets were activated with ADP and epinephrine (10  $\mu$ M each). The data shown are from a single experiment representative of two runs performed as described.

Figure 5 shows the effect of Arg-Tyr-Asp-21mer on platelet aggregation. Gel-filtered platelets were stirred for 3 min in an aggregometer cuvette in the presence of ADP (10  $\mu$ M), fibrinogen (100  $\mu$ g/ml),  $\text{CaCl}_2$  (1mM), and one of the following: Arg-Tyr-Asp-21mer (filled circles); Arg-Tyr-Asp-21mer having a Gly substituted for the Tyr residue

(Arg-Gly-Asp-21mer) (filled squares); Arg-Tyr-Asp-21mer having the Asp and Arg residues reversed (Asp-Tyr-Arg-21mer) (filled triangles); or Arg-Tyr-Asp-21mer having a d-Ala residue substituted for the Tyr residue (Arg-d-Ala-Asp-21mer) (unfilled circles). Aggregation is expressed as the initial rate of aggregation, where the rate of aggregation in the absence of added peptide is arbitrarily denoted as 100%. The data represent the means of three separate experiments performed as described.

Figure 6 shows the effect of Arg-Tyr-Asp on the binding of FITC-PAC1 (Figure 6A) or FITC-9F9 (Figure 6B) and fibrinogen to platelets stimulated with ADP and epinephrine (10  $\mu$ M each). The specific peptides tested were Arg-Tyr-Asp-21mer (filled circles), Arg-Gly-Asp-21mer (filled squares), Asp-Tyr-Arg-21mer (filled triangles), Arg-d-Ala-Asp-21mer (unfilled circles) or Arg-Gly-Asp-Ser (unfilled squares). The conditions in each of the binding assays were identical to those in Figure 3. Data shown are from a single experiment representative of three runs performed as described.


#### DETAILED DESCRIPTION OF THE INVENTION

The present invention involves the discovery of novel and very active compounds that contain the tripeptide sequence of arginine-tyrosine-aspartic acid, linked from amino- to carboxy-terminus. The subject invention derives from the surprising finding that the arginine-tyrosine-aspartic acid sequence is present in one of the heavy chain hypervariable regions of PAC1 murine monoclonal immunoglobulin, and in that position is a potential mediator in the binding of this monoclonal antibody to the blood platelet membrane fibrinogen receptor, glycoprotein IIb-IIIa.

More specifically, the present invention encompasses peptide compounds of the general Formula I



and physiologically acceptable salts thereof. The present invention further encompasses peptide compounds of the general Formula II



and physiologically acceptable salts thereof. These compounds are characterized by their inhibitory potency, namely their ability to retard or prevent the following: the binding of the adhesive protein fibrinogen to blood platelets (referred to herein as inhibiting "fibrinogen binding"); the binding blood platelets to themselves (referred to herein as inhibiting "platelet aggregation"); and/or the binding of compounds or substances, particularly proteins, to the glycoprotein IIb-IIIa complex found in blood platelet membranes (referred to herein as inhibiting "glycoprotein IIb-IIIa binding"). The subject com-

pounds are also useful in retarding or preventing the formation of blood clots or thrombi (referred to herein as inhibiting "thrombosis") and/or the spread of cancer cells throughout the body (referred to herein as inhibiting "cancer metastasis"). The selectivity of these compounds in carrying out the foregoing related activities makes them particularly useful as therapeutic and/or diagnostic agents.

As used throughout this specification, amino acid residues are denoted by the following abbreviations:

Alanine = Ala

Arginine = Arg

Asparagine = Asn

Aspartic acid = Asp

Cysteine = Cys

Glutamic acid = Glu

Glutamine = Gln

Glycine = Gly

Histidine = His

Isoleucine = Ile

Leucine = Leu

Lysine = Lys

Methionine = Met

Phenylalanine = Phe

Proline = Pro

Serine = Ser

Threonine = Thr

Tryptophan = Trp

Tyrosine = Tyr

Valine = Val

For maximal activity, the amino acid residues comprising the key tripeptide region, Arg-Tyr-Asp, of the subject invention must be in the L-configuration. The remaining amino acid residues of the subject invention can be present in either the D- or the L-configuration.

As previously noted, in the above Formula I,  $X^1$  and  $X^2$ , and in the above Formula II,  $X^1$ ,  $X^2$  and Y, may each independently comprise one or more amino acid residues. The  $X^1$ ,  $X^2$  and/or Y amino acid residues, as the case may be, may be linked among themselves, or with each other, in branched, cyclical or straight chain form, although straight chain linkages are preferred. As one skilled in the art would recognize, branched or cyclical chains may be produced by the formation of a peptide bond with amino acid side groups that contain amino or carboxyl moieties. Amino acids containing such side groups include, for example, glutamic acid (carboxyl group), aspartic acid (carboxyl group) and lysine (amide group). Branched or cyclical chains may also be produced through the formation of a covalent disulfide bond between amino acid residues having sulfur-containing side groups, such as cysteine.





In Formula I, as noted above, m and n may be,

independently, 0 or 1 -- that is, m and n may both be 0 or may both be 1, or m may be 0 when n is 1, or m may be 1 when n is 0. Where both m and n are 0, the subject invention comprises a tripeptide of the sequence Arg-Tyr-Asp, linked, in order, from amino- to carboxy-terminus. Similarly, in Formula II, when each of m, n and o is 0, the subject invention is also directed to the tripeptide Arg-Tyr-Asp.

The R<sup>1</sup> substituent of Formulas I or II may be either an unprotected or protected terminal amino group (amino-terminus). Similarly, R<sup>2</sup> may be either an unprotected or protected terminal carboxy group (carboxy-terminus).

As used herein, "protected" terminal amino group, refers to a terminal amino group coupled with any of various amino-terminal protecting groups traditionally employed in peptide synthesis. Examples of suitable groups include acyl protecting groups, for example, formyl, acetyl, benzoyl, trifluoroacetyl, succinyl and methoxysuccinyl; aromatic urethane protecting groups, for example, benzyloxycarbonyl; and aliphatic urethane protecting groups, for example, tert-butoxycarbonyl or adamantyloxycarbonyl. Gross and Mienhofer, eds., *The Peptides*, Vol. 3, pp. 3-88 (Academic Press, New York, 1981), disclose numerous suitable terminal amino protecting groups.

The following represent preferred amino terminal protecting groups:



The amino acid residues of the X<sup>1</sup>, X<sup>2</sup> and Y substituents having side-chain amino groups, for

example, Lys or Arg, can optionally comprise suitable amino-terminal protecting groups attached to the side chains.

As used herein, "protected" terminal carboxyl group, refers to a terminal carboxyl group coupled with any of various carboxy-terminal protecting groups. As will be readily apparent to one skilled in the art, suitable groups include tert-butyl, benzyl or other acceptable groups linked to the terminal carboxyl group through an ester or ether bond. Amino acid residues of the X<sup>1</sup>, X<sup>2</sup> and Y substituents having acidic or hydroxy side chains can be similarly protected.

The present invention also contemplates physiologically acceptable salts of Formulas I and II. These salts include acid addition salts, for example, salts of hydrochloric acid, hydrobromic acid, acetic acid, trifluoroacetic acid, citric acid, succinic acid, benzene sulfonic acid or other suitable acid addition salts.

Preferable classes of compounds within the scope of Formula I include:

o Compounds wherein

m is 0; and

n is 0.

o Compounds wherein X<sup>1</sup> and X<sup>2</sup> are in straight chain forms.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup> and X<sup>2</sup>, taken together, is less than about 40.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup> and X<sup>2</sup>, taken together, is less than about 25.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup> and X<sup>2</sup>, taken together, is less than about 10.

o Compounds wherein

X<sup>1</sup> is Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr;

m is 1; and

n is 1.

o Compounds wherein

X<sup>1</sup> is Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr;

m is 1; and

n is 1.

o Compounds wherein

X<sup>1</sup> is Gln-Val-Gln-Leu-Lys-Gln-Ser-Gly-Pro Gly-Leu-

Val-Gln-Pro-Ser-Gln-Ser-Leu-Ser-Ile-Thr-Cys-Thr-

Val-Ser-Gly-Phe-Ser-Leu-Thr-Ser-Tyr-Gly-Val-His-

Trp-Val-Arg-Leu-Ser-Pro-Gly-Lys-Gly-Leu-Glu-Trp-

Leu-Gly-Val-Ile-Trp-Ser-Gly-Gly-Ser-Thr-Asp-Tyr-

Asn-Ala-Ala-Phe-Ile-Ser-Arg-Leu-Ser-Ile-Ser-Lys-

Asp-Asn-Ser-Lys-Ser-Gln-Val-Phe-Phe-Lys-Met-

Asn-Ser-Leu-Gln-Ala-Asn-Asp-Thr-Gly-Ile-Tyr-Try-

Cys-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp Tyr-Trp-

Gly-Gln-Gly-Thr-Ser-Val-Thr-Val-Ser-Ser;

m is 1; and

n is 1.

o Compounds wherein

X<sup>1</sup> is Gln-Val-Gln-Leu-Lys-Gln-Ser-Gly-Pro Gly-Leu-Val-Gln-Pro-Ser-Gln-Ser-Leu-Ser-Ile-Thr-Cys-Thr-Val-Ser-Gly-Phe-Ser-Leu-Thr-Ser-Tyr-Gly-Val-His-Trp-Val-Arg-Leu-Ser-Pro-Gly-Lys-Gly-Leu-Glu-Trp-Leu-Gly-Val-Ile-Trp-Ser-Gly-Gly-Ser-Thr-Asp-Tyr-Asn-Ala-Ala-Phe-Ile-Ser-Arg-Leu-Ser-Ile-Ser-Lys-Asp-Asn-Ser-Lys-Ser-Gln-Val-Phe-Phe-Lys-Met-Asn-Ser-Leu-Gln-Ala-Asn-Asp-Thr-Gly-Ile-Tyr-Tyr-Cys-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp Tyr-Trp-Gly-Gln-Gly-Thr-Ser-Val-Thr-Val-Ser-Ser;

m is 1; and

n is 1.

o Compounds wherein

R<sup>1</sup> is an unprotected terminal amino group; and

R<sup>2</sup> is an unprotected terminal carboxy group.

Preferable classes of compounds within the scope of Formula II include:

o Compounds wherein

m is 0;

n is 0; and

o is 0. o Compounds wherein X<sup>1</sup>, and X<sup>2</sup> and Y are in straight chain forms.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, and X<sup>2</sup> and Y, taken together, is less than about 250.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, and X<sup>2</sup> and Y, taken together, is less than about 200.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, and X<sup>2</sup> and Y, taken together, is less than about 100.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, X<sup>2</sup> and Y, taken together, is less than about 50.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, X<sup>2</sup> and Y, taken together, is less than about 40.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, X<sup>2</sup> and Y, taken together, is less than about 25.

o Compounds wherein the total number of amino acid residues in X<sup>1</sup>, X<sup>2</sup> and Y, taken together, is less than about 10.

o Compounds wherein

X<sup>1</sup> is Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr;

m is 1;

n is 1; and

o is 0.

o Compounds wherein

X<sup>1</sup> is Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr;

m is 1;

n is 1; and

o is 0.

o Compounds wherein

X<sup>1</sup> is Gln-Val-Gln-Leu-Lys-Gln-Ser-Gly-Pro Gly-Leu-Val-Gln-Pro-Ser-Gln-Ser-Leu-Ser-Ile-Thr-Cys-Thr-Val-Ser-Gly-Phe-Ser-Leu-Thr-Ser-Tyr-Gly-Val-His-Trp-Val-Arg-Leu-Ser-Pro-Gly-Lys-Gly-Leu-Glu-Trp-Leu-Gly-Val-Ile-Trp-Ser-Gly-Gly-Ser-Thr-Asp-Tyr-Asn-Ala-Ala-Phe-Ile-Ser-Arg-Leu-Ser-Ile-Ser-Lys-Asp-Asn-Ser-Lys-Ser-Gln-Val-Phe-Phe-Lys-Met-Asn-Ser-Leu-Gln-Ala-Asn-Asp-Thr-Gly-Ile-Tyr-Tyr-Cys-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp Tyr-Trp-Gly-Gln-Gly-Thr-Ser-Val-Thr-Val-Ser-Ser;

m is 1;

15 n is 1; and

o is 0.

o Compounds wherein

X<sup>1</sup> is Gln-Val-Gln-Leu-Lys-Gln-Ser-Gly-Pro Gly-Leu-Val-Gln-Pro-Ser-Gln-Ser-Leu-Ser-Ile-Thr-Cys-Thr-Val-Ser-Gly-Phe-Ser-Leu-Thr-Ser-Tyr-Gly-Val-His-Trp-Val-Arg-Leu-Ser-Pro-Gly-Lys-Gly-Leu-Glu-Trp-Leu-Gly-Val-Ile-Trp-Ser-Gly-Gly-Ser-Thr-Asp-Tyr-Asn-Ala-Ala-Phe-Ile-Ser-Arg-Leu-Ser-Ile-Ser-Lys-Asp-Asn-Ser-Lys-Ser-Gln-Val-Phe-Phe-Lys-Met-Asn-Ser-Leu-Gln-Ala-Asn-Asp-Thr-Gly-Ile-Tyr-Tyr-Cys-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;

X<sup>2</sup> is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp Tyr-Trp-Gly-Gln-Gly-Thr-Ser-Val-Thr-Val-Ser-Ser;

m is 1;

30 n is 1; and

o is 0.

o Compounds wherein

R<sup>1</sup> is an unprotected terminal amino group; and

R<sup>2</sup> is an unprotected terminal carboxy group.

35 Obvious equivalents of the foregoing compounds include compounds comprising less common or modified amino acids, for example, hydroxyproline, hydroxylysine, cystine, thyroxine, norleucine, pyroglutamic acid or other derivatives capable of incorporation into the peptides of the present invention.

The branched, cyclical and straight chain peptides of Formulas I and II can be synthesized using any one of a number of conventional preparative and recovery methods, as will be readily apparent to one skilled in the art.

45 A preferable synthesis route for the straight chain, especially the smaller, peptides of the invention is the solid phase method. This method is well known in the art and is described, for example, in U.S. Patent Nos. 4,683,291, 4,244,946, 4,305,872, and 4,316,891, and in Merrifield, *J. Am. Chem. Soc.*, 85 2149 (1964), Vale et al, *Science* 213, 1394-1397 (1981) and Marke et al, *J. Am. Chem. Sci.*, 103, 3178 (1981), the disclosures of which are incorporated herein by reference. Other preparative methods which may be employed include the processes of Houghten, *Proc. Natl. Acad. Sci.*, 82,

5132 (1985), the disclosures of which are also incorporated herein by reference. A preferable synthesis procedure particularly for the smaller branched or cyclical chain peptides, as a skilled artisan would recognize, would include conventional liquid phase processes. The liquid phase processes, as well as other synthesis methods, are described in Principles of Peptide Synthesis, M. Bodansky, ed. (Springer-Verlag, 1984), the disclosures of which are incorporated herein by reference. As would be apparent to one cognizant of the art, for larger peptides, a preferable procedure may include conventional recombinant DNA techniques. Such techniques are discussed in greater detail below. Recombinant techniques as well known and are described, for example, in Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor (1982), the disclosures of which are incorporated herein by reference.

Suitable recovery methods are described in the foregoing publications. Other recovery methods which may be employed include those described in Rivier et al., Peptides: The Structure and Biological Function, pp. 125-128 (1979), the disclosures of which are incorporated herein by reference.

The invention further provides compositions and methods for utilizing the subject compounds and compositions in inhibiting fibrinogen binding, platelet aggregation, and glycoprotein IIb-IIIa binding. The ability of the subject compounds to inhibit the foregoing activities makes them useful in inhibiting the physiological process of thrombosis. In addition, in light of their demonstrated activities, the compounds of the present invention may be employed in inhibiting cancer metastasis, an aberrant physiological phenomenon that is believed to require the adhesion of blood platelets to the cancer cells. The specificity of the subject compounds in carrying out these related functions makes them particularly useful as therapeutic and/or diagnostic agents.

In particular, the compositions of the present invention comprise an effective amount of a compound of Formula I and a physiologically acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic, diagnostic or other uses are well-known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Gennaro, A.R., ed. (Mack Publishing Co., Easton PA, 1985).

In practicing the methods of the invention, the compounds or compositions can be used alone or in combination with one another, or in combination with other therapeutic or diagnostic agents. They can be utilized in vivo, ordinarily in a mammal, preferably in a human, or in vitro. In employing in vitro, the compounds or compositions are applied to a cellular solution, that is, a solution containing

the cellular component or components sought to be inhibited. In employing in vivo, the compounds or compositions can be administered to the mammal in a variety of ways, including parenterally, intravenously, subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety of dosage forms. As will be readily apparent to one skilled in the art, the useful in vivo dosage to be administered and the particular mode of administration will vary depending upon the age, weight and mammalian species treated, the particular compounds employed, and the specific use for which these compounds are employed. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be well within the ambit of one skilled in the art. Typically, applications are commenced at lower dosage levels, with dosage being increased until the desired effect is achieved.

Antibodies, both monoclonal and polyclonal, directed to peptide compounds of the present invention are useful in isolation and identification of the subject proteins, and the present invention also pertains to such antibodies. To prepare the subject antibodies, any one of a number of techniques which known in the art can be employed. In one such technique, polyclonal antibodies may be synthesized by injecting an animal (for example, a rabbit) with one or more compounds of the invention. After injection, the animal naturally produces antibodies to these compounds. When the antibody level rises to a sufficient level, antibody-containing blood, called antiserum, is then drawn from the animal, and the compound-specific antibody is isolated from other antibodies in the antiserum by any one of a number of separation techniques (for example, affinity chromatography). Monoclonal antibodies may be prepared using the technique of Kohler and Milstein, Nature 256, pp. 495-497 (1975).

The invention further relates to nucleic acid, preferably DNA, sequences which encode the peptide compounds of the present invention. Such fragments may be synthesized using standard nucleic acid synthesis techniques or, alternatively, may be obtained from genomic material of various sources using conventional genetic engineering protocol. One source of genomic material is the PAC1 hybridoma cell described in Shatill et al., J. Biol. Chem., 260, 11107-11114 (1985), particularly the PAC1- $\mu$  heavy chain variable region of that genome. Using standard recombinant DNA techniques, the nucleic acid sequences can be, if desired, incorporated into an appropriate cloning vector, transduced into an appropriate host cell and the proteins encoded by the nucleic acid fragment expressed and collected for further use, thereby providing a potentially convenient method for the



production of the proteins, particularly the lengthy straight chain proteins, of the subject invention. Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor (1982), provides a detailed discussion of various recombinant DNA methodologies.

Cloning vector as used herein is defined as a DNA (or RNA) sequence, which allows the incorporation of specific experimental foreign DNA (or RNA), with the combined DNA (or RNA) being introduced into a host cell that can exist in a stable manner and express the protein dictated by the experimental DNA (or RNA). The foreign DNA (or RNA) combined with the vector DNA (or RNA) constitutes a recombinant DNA (or RNA) molecule which is derived from recombinant technology. Cloning vectors may include plasmids, bacteriophage, viruses and cosmids. Expression vectors are defined herein as DNA (or RNA) sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express either procaryotic or eucaryotic genes in a variety of cells such as bacteria, yeast, insect and mammalian cells. The proteins may also be expressed in a number of virus systems. An appropriately constructed expression vector should contain an origin of replication for autonomous replication in host cells, selective markers, a limited number of useful restriction enzyme sites, a high copy number, and strong promoters. A promoter is defined as a DNA (or RNA) sequence that directs RNA polymerase to bind to DNA and to initiate RNA synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

Finally, the present invention includes a peptide comprising an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\mu$  heavy chain of monoclonal antibody PAC1, as shown in Figure 2, and any antibodies incorporating such sequence. The present invention further includes a peptide comprising an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\mu$  heavy chain of monoclonal antibody PAC1, as shown in Figure 2, and an amino acid sequence corresponding to the amino acid sequence of the variable region of the  $\kappa$  light chain of monoclonal antibody PAC1, as shown in Figure 1, and any antibodies incorporating such sequences. As will be apparent to those skilled in the art, and as described in the foregoing discussion of peptide synthesis using chemical and recombinant methodologies such as described in Riechmann et al., Nature, Vol. 332, pp. 323-327 (1988), various tech-

niques are available to prepare such peptides. The recombinant DNA techniques can also be applied in the area of antibody manipulation. For a discussion of antibody engineering, see Riechmann et al., Nature, 332, 323-327 (1988), the disclosures of which are herein incorporated by reference.

Most preferably, the invention is directed to monoclonal antibody PAC1. The hybridoma cell line producing this monoclonal antibody is described in Shattil et al., J. Biol. Chem., 260, 11107-11114 (1985), the disclosures of which are hereby incorporated by reference. Both the hybridoma cell PAC1 and the PAC1 monoclonal antibody produced therefrom are discussed in the Examples below. As with the peptides of Formulas I and II, the peptides and monoclonal antibodies of this paragraph are useful as inhibitors of fibrinogen binding, platelet aggregation, and/or glycoprotein IIb-IIIa binding, and, similarly, are effective in the treatment of certain physiological conditions, such as thrombosis and/or cancer metastasis, and the present invention is further directed to these uses.

The invention is further described in the following Examples. These Examples are not to be construed as limiting the scope of the appended Claims.

## Examples

### Example 1

#### Cell Culture and Isolation of mRNA

PAC1 hybridoma cells, prepared as described in Shattil et al., J. Biol. Chem., 260, 11107-11114 (1985), were grown in a standard culture medium containing fetal calf serum, and  $5 \times 10^8$  cells were harvested and resuspended in 4M guanidine thiocyanate as described in Chirgwin et al., Biochemistry, 18, 5294-5299 (1979). Total RNA was isolated after centrifugation through a CsCl cushion and poly A+ RNA was isolated using oligodT cellulose following the procedures set forth in Aviv et al., Proc. Natl. Acad. Sci., U.S.A., 69, 264-268 (1972).

### Example 2

#### Sequencing of PAC1 Immunoglobulin Variable Regions

Using complementary oligonucleotide primers from the 5' region of the mouse  $\mu$  and  $\kappa$  constant



region mRNAs, (5'-GCTCTCGCAGGAGACGAG and 5'-GGTGGGAAGATGGATACAGTT, respectively), the sequences of PAC1- $\mu$  (light chain) and  $\kappa$  (heavy chain) mRNAs were determined using the direct dideoxy chain termination method as described by Geliebter in *Focus*, 9, 1,5-8 (1987). Briefly, 20 ng of end-labeled primers were annealed to 10  $\mu$ g of poly A+ mRNA at 50°C and primers were extended in the presence of dideoxynucleotides and reverse transcriptase. The products of this reaction were electrophoresed on 6% denaturing polyacrylamide gels that were then exposed to autoradiography. Once approximately 200 bp of sequence had been determined, complementary primers within the variable region about 150 bp upstream of the constant region were synthesized in order to extend the sequence further 5' within the variable region. The primer used in the PAC1- $\mu$  reaction was 5'-CATTAGCTTG-CAGACTGTTC and in the PAC1- $\kappa$  reaction was 5'-ATCTTGAGTGTGAAATCTGT. A small portion, about 50 nucleotides, of the 3' end of the  $\kappa$  mRNA was determined using the modified Maxam and Gilbert technique of Schlomchik et al., *J. Exp. Med.*, 164, 407-427 (1986). Figure 1 shows the nucleotide sequence of the variable region of the  $\kappa$  light chain of a PAC1 hybridoma cell, along with the predicted amino acid sequence. Figure 2 shows the nucleotide sequence of the variable region of the  $\mu$  heavy chain of a PAC1 hybridoma cell, along with the predicted amino acid sequence. In both Figures 1 and 2, the complementary determining regions (CDRs) are shown in boxes. The amino acids numbered according to convention. Kabat E.A., Wu, T.T., Reid-Miller, M., Perry, H.M., Gottesman, K.S., *Sequences of Proteins of Immunologic Interest* (US DHHS, PHS, NIH 1987).

#### Example 3

##### Peptide Synthesis and Analysis

The peptide NH<sub>2</sub>-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr-Arg-Tyr-Asp-Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr-COOH, which corresponds to amino acids 93 through 102 of the PAC1- $\mu$  variable region (Figure 2) and which is referred to herein as the Arg-Tyr-Asp-21mer, and various other peptides used herein were synthesized by the solid phase method using an Applied Biosystems Inc. 430A automated synthesizer. Boc-amino acids supplied by the manufacturer were used except for Boc-Asp (Chx). Triple couplings were used for Arg<sup>9</sup>, Ser<sup>4</sup>, Tyr<sup>7</sup> Tyr<sup>8</sup>, Arg<sup>3</sup>, Arg<sup>1</sup> and Ala<sup>1</sup> of the Arg-Tyr-Asp-21mer. The remaining residues were incorporated by a double coupling protocol. The peptides were

cleaved from the solid support by the Sn2/Sn1 HF cleavage procedure described in Tam, *J. Amer. Chem. Soc.*, 105, 6442-6455 (1983) and purified by gel filtration (Sephadex G-50F, 50% HOAc) and reverse phase HPLC (Vydac C18, 15 $\mu$ , 5 x 30 cm column, 95% A-B to 75% A-B over 60 minutes at a flow rate of 100 ml/min; A = 0.1% TFA-H<sub>2</sub>O, B = 0.1% TFA-CH<sub>3</sub>CN). Products were characterized by amino acid analyses after acid hydrolysis, fast atom bombardment mass spectrometry (FABMS), nuclear magnetic resonance spectroscopy, and sequence analysis after Edman degradation. Analyses were consistent with the expected structures, with a product purity of  $\geq$  95%.

#### Example 4

##### Binding of PAC1 and Fibrinogen to Platelets

The binding of PAC1 monoclonal antibody and fibrinogen to gel-filtered human blood platelets was examined by fluorescence-activated flow cytometry using the procedures of Shattil et al., *Blood*, 70, 307-315 (1987). Venous blood from normal donors was drawn into 1/7 vol of NIH formula A acid-citrate-dextrose solution, platelet-rich plasma was obtained, and the platelets were gel-filtered into an isotonic HEPES-containing buffer at pH 7.4, as described in Shattil et al., *Blood*, 68, 1224-1231 (1986). To examine the effect of synthetic peptides on PAC1 binding to activated platelets, 5 x 10<sup>7</sup> platelets were incubated for 15 min at room temperature in a final vol of 50  $\mu$ l in the presence of varying concentrations of (0-100  $\mu$ M) of peptide, 40  $\mu$ g/ml fluorescein-labeled PAC1 (FITC-PAC1), 10  $\mu$ M ADP and 10  $\mu$ M epinephrine. Then, 500  $\mu$ l of the isotonic buffer was added and antibody binding was quantitated using a flow cytometer, as described in Shattil et al., *Blood*, 70, 307-315 (1987). Fibrinogen binding to platelets was measured using the same incubation system, except that 50  $\mu$ g/ml of purified human fibrinogen as described in Bennett et al., *J. Clin. Invest.*, 64, 1393-1400 (1979) was added and the monoclonal antibody, FITC-9F9, was used instead of FITC-PAC1. Antibody 9F9 is specific for fibrinogen and has been used successfully to measure the amount of fibrinogen bound to platelets. Abrams et al., *Blood*, 70, 335a (1987). Previous studies have established that the amount of FITC-labeled antibody bound to platelets determined by flow cytometry is related in a linear fashion to the amount of antibody bound determined in a conventional radioligand binding assay. Shattil et al., *Blood*, 70, 307-315 (1987). The amount of platelet-bound FITC-PAC1 or FITC-9F9 was determined by analyzing 10,000 platelets for

the extent of fluorescence at 488 nm, and binding expressed as the mean fluorescence intensity in arbitrary fluorescence units.

Under these conditions, the Arg-Tyr-Asp-21mer peptide inhibited the binding of FITC-PAC1 to platelets with an  $IC_{50}$  of 20-30  $\mu$ M (as shown in Figure 3A, filled circles). This was similar to the  $IC_{50}$  of the tetrapeptide,  $NH_2$ -Arg-Gly-Asp-Ser-COOH (Arg-Gly-Asp-Ser) (Figure 3A, unfilled circle). In five separate experiments, Dixon plot analysis carried out in accordance with Segel, Enzyme Kinetics 465-504 Wiley-Interscience, N.Y. (1975), revealed that the apparent  $K_i$  for inhibition of PAC1 binding by Arg-Tyr-Asp-21mer was  $10.2 \pm 4.3$   $\mu$ M (SE). As a control, an unrelated peptide of similar length having the sequence  $NH_2$ -Tyr-Val-Asp-Gly-Asp-Gln-Cys-(Acm)-Glu-Ser-Asn-Pro-Cys-Leu-Asn-Gly-Gly-Met(O)-Cys-(Acm)-Lys-Asp-Asp-Ile-Asn-Ser-Tyr-Gly-Cys-Phe-COOH (unrelated-28mer) (control) had no effect on FITC-PAC1 binding (Figure 3A, unfilled squares). Identical results were obtained when platelets were activated with phorbol myristate acetate instead of ADP and epinephrine (data not shown).

The ability of Arg-Tyr-Asp-21mer to inhibit fibrinogen binding to activated platelets was also tested. Fibrinogen binding was assessed by flow cytometry using a FITC-labeled anti-fibrinogen monoclonal antibody, 9F9. When platelets were activated with ADP plus epinephrine in the presence of 50  $\mu$ g/ml of fibrinogen, Arg-Tyr-Asp-21mer inhibited fibrinogen binding with an  $IC_{50}$  of  $\mu$ M (Figure 3B, filled circle). The control peptide (unrelated 28 mer) had no effect on 9F9 binding (Figure 3B, unfilled square). In three experiments, Dixon plot analysis showed that this peptide inhibited fibrinogen binding with an apparent  $K_i$  of  $5.5 \pm 3.5$   $\mu$ M.

To examine the specificity of the Arg-Tyr-Asp-21mer peptide in inhibiting PAC1 monoclonal antibody binding, the effect of this peptide on the binding of monoclonal antibodies to other platelet glycoproteins were examined. The Arg-Tyr-Asp-21mer peptide had no effect on the binding of FITC-AP1 to platelet glycoprotein Ib on resting platelets (Figure 4, unfilled circles). In addition, the peptide had no effect on the binding of either FITC-B1B5 to glycoprotein IIb or of FITC-S12 to GMP-140 on activated platelets (Figure 4, filled squares and unfilled triangles, respectively). The peptide had a marked effect on PAC1 binding (Figure 4, filled circles). Antibody S12, described in McEver et al., J. Biol. Chem., 259: 9799 (1984) specific for an  $\alpha$ -granule membrane protein (GMP) expressed on the surface of activated platelets. Antibody B1B5 is specific for glycoprotein IIb, and is discussed in Shattil, J. Biol. Chem., 260, 11107-11114 (1985) and Bennett et al., Proc. Natl. Acad. Sci.,

U.S.A., 80, 2417-2421 (1983).

#### Example 5

#### Platelet Aggregation Studies

The Effect on Arg-Tyr-Asp-21mer peptide on blood platelet aggregation was examined. The aggregation of gel-filtered platelets was carried out as described in Shattil et al., Blood, 68, 1224-1231 (1986). Platelets ( $2 \times 10^8$ /ml) were stirred at 37°C in the presence of human fibrinogen (100  $\mu$ g/ml) and  $CaCl_2$  (1 mM) and aggregation was initiated by the addition of 10  $\mu$ M ADP. Aggregation was monitored as a change in light transmittance, and is expressed as the initial rate of aggregation.

In three experiments Arg-Tyr-Asp-21mer inhibited the initial rate of ADP-induced platelet aggregation with an  $IC_{50}$  of approximately 40  $\mu$ M (Figure 5, filled circles). At 10  $\mu$ g/ml fibrinogen the  $IC_{50}$  was approximately 10  $\mu$ M (data not shown).

#### Example 6

#### Modifications in the Arg-Tyr-Asp Sequence of the 21mer Peptide

The specific role of the Arg-Tyr-Asp sequence within Arg-Tyr-Asp-21mer was examined by testing three additional peptides identical to the Arg-Tyr-Asp-21mer except for specific modifications in the Arg-Tyr-Asp region. Substitution of the tyrosine in Arg-Tyr-Asp with glycine to produce an Arg-Gly-Asp-21mer increased the inhibitory potency of the peptide by about ten-fold in both the aggregation assay (Arg-Gly-Asp-21mer, Figure 5, filled squares; Arg-Tyr-Asp-21mer, Figure 5, filled circles) and the PAC1 and fibrinogen binding assays (Arg-Tyr-Asp-21mer, Figure 6, filled squares; Arg-Tyr-Asp-21mer, Figure 6, unfilled squares). On the other hand, substitution of the tyrosine with a D-alanine to produce an Arg-d-Ala-Asp-21mer, or inversion of the Arg and Asp residues to produce an Arg-Tyr-Asp-21mer substantially reduced the inhibitory capacity of the peptide (Arg-d-Ala-Asp-21mer, Figure 6, unfilled circles; Asp-Tyr-Arg-21mer, Figure 6, filled triangles). The Arg-Gly-Asp-Ser-tetramer is shown in Figure 6 as the unfilled square.

These specific modifications of the Arg-Tyr-Asp sequence within the Arg-Tyr-Asp-21mer peptide clearly establish that this sequence of three amino acids is of significance to the functional properties of the entire peptide.

Various modifications of the invention in addi-

tion to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended Claims.

### Claims

1. A compound of the formula  
 $R^1-(X^1)_n-Arg-Tyr-Asp-(X^2)_m-R^2$   
 and physiologically acceptable salts thereof,  
 wherein:  
 $X^1$  and  $X^2$ , independently, are 1 to about 50 amino  
 acid residues, provided that the total number of  
 amino acid residues in  $X^1$  and  $X^2$ , taken together, is  
 equal to or less than about 50;  
 $m$  and  $n$ , independently, are 0 or 1;  
 $R^1$  is a protected or unprotected terminal amino  
 group; and  
 $R^2$  is a protected or unprotected terminal carboxy  
 group.
2. A compound according to Claim 1 wherein  
 $m$  is 0; and  
 $n$  is 0.
3. A compound according to either of Claims 1  
 or 2 wherein  $X^1$  and  $X^2$  are in straight chain form.
4. A compound according to any preceding  
 Claim wherein the total number of amino acid resi-  
 dues in  $X^1$  and  $X^2$ , taken together, is less than  
 about 40.
5. A compound according to any of Claims 1 to  
 3 wherein the total number of amino acid residues  
 in  $X^1$  and  $X^2$ , taken together, is less than about 25.
6. A compound according to any of Claims 1 to  
 3 wherein the total number of amino acid residues  
 in  $X^1$  and  $X^2$ , taken together, is less than about 10.
7. A compound according to Claim 1 wherein  
 $X^1$  is Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;  
 $X^2$  is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr;  
 $m$  is 1; and  
 $n$  is 1.
8. A compound according to Claim 1 wherein  
 $X^1$  is Gln-Val-Gln-Leu-Lys-Gln-Ser-Gly-Pro-Gly-Leu-  
 Val-Gln-Pro-Ser-Gln-Ser-Leu-Ser-Ile-Thr-Cys-Thr-  
 Val-Ser-Gly-Phe-Ser-Leu-Thr-Ser-Tyr-Gly-Val-His-  
 Trp-Val-Arg-Leu-Ser-Pro-Gly-Lys-Gly-Leu-Glu-Trp-  
 Leu-Gly-Val-Ile-Trp-Ser-Gly-Gly-Ser-Thr-Asp-Tyr-  
 Asn-Ala-Ala-Phe-Ile-Ser-Arg-Leu-Ser-Ile-Ser-Lys-  
 Asp-Asn-Ser-Lys-Ser-Gln-Val-Phe-Phe-Lys-Met-  
 Asn-Ser-Leu-Gln-Ala-Asn-Asp-Thr-Gly-Ile-Tyr-Tyr-  
 Cys-Ala-Arg-Arg-Ser-Pro-Ser-Tyr-Tyr;  
 $X^2$  is Gly-Ala-Gly-Pro-Tyr-Tyr-Ala-Met-Asp-Tyr-Trp-  
 Gly-Gln-Gly-Thr-Ser-Val-Thr-Val-Ser-Ser;  
 $m$  is 1; and  
 $n$  is 1.
9. A compound of the formula  
 $R^1-(X^1)_n-Arg-Tyr-Asp-(Y)_o(X^2)_m-R^2$   
 and physiologically acceptable salts thereof,  
 wherein:  
 $X^1$  and  $X^2$ , independently, are 1 or more amino  
 acid residues;  
 $Y$  is 1 or more amino acid residues, provided that  $Y$   
 does not include a Ser or Ala residue;  
 $m$ ,  $n$  and  $o$ , independently, are 0 or 1, provided  
 that when  $m$  is 1,  $o$  must be 1  
 $R^1$  is a protected or unprotected terminal amino  
 group; and  
 $R^2$  is a protected or unprotected terminal carboxy  
 group; and wherein said compound inhibits  
 fibrinogen binding, platelet aggregation, thrombo-  
 sis, cancer metastasis and/or glycoprotein IIb-IIIa  
 binding.
10. A composition for inhibiting, fibrinogen  
 binding, platelet aggregation, thrombosis, and/or  
 cancer metastasis comprising an effective amount  
 of a compound of any preceding Claim and a  
 physiologically acceptable carrier or diluent there-  
 for.
11. A composition for inhibiting binding of a  
 substance to a blood platelet membrane  
 glycoprotein IIb-IIIa complex comprising an effec-  
 tive amount of a compound of any of Claims 1 to 9  
 and a physiologically acceptable carrier diluent.
12. A method of inhibiting fibrinogen binding,  
 platelet aggregation, comprising administering to  
 the platelets an effective amount of a compound of  
 any of Claims 1 to 9.
13. A method of inhibiting binding of a sub-  
 stance to a blood platelet membrane glycoprotein  
 IIb-IIIa complex comprising administering to the  
 complex an effective amount of a compound of any  
 of Claims 1 to 9.
14. An antibody to a compound of Claim 1.
15. A nucleic acid fragment comprising a  
 nucleotide sequence which encodes a compound  
 of any of Claims 1 to 9.
16. A peptide comprising an amino acid se-  
 quence corresponding to the amino acid sequence  
 shown in Figure 2.
17. A peptide according to Claim 16 further  
 comprising an amino acid sequence corresponding  
 to the amino acid sequence as shown in Figure 1.
18. A monoclonal antibody comprising an ami-  
 no acid sequence corresponding to the amino acid  
 sequence shown in Figure 2.
19. A monoclonal antibody according to Claim  
 18, further comprising an amino acid sequence  
 corresponding to the amino acid sequence shown  
 in Figure 1.
20. The use in the manufacture of a medica-  
 ment effective *in vivo* for inhibiting fibrinogen bind-  
 ing, platelet aggregation, thrombosis, and/or cancer  
 metastasis which comprises a composition accord-  
 ing to any of Claims 1 to 9




**PAC1- $\mu$  Variable Region**

[illegible]

**FIGURE 3**

Effect of Arg-Tyr-Asp-21mer on Binding of  
PAC1 and Fibrinogen to Activated Platelets



**FIGURE 4**

**Effect of Arg-Tyr-Asp-2imer on Binding of  
Monoclonal Antibodies to Platelets**






FIGURE 5

Effect of Arg-Tyr-Asp-dimer on  
ADP-Induced Platelet Aggregation





FIGURE 6

Effect of Arg-Tyr-Asp-21mer on the Binding of  
PAC1 and 9F9 to Activated Platelets



(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 368 486 A3**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 89310620.3

(22) Date of filing: 17.10.89

(51) Int. Cl.<sup>5</sup>: C07K 5/08, C07K 7/06,  
C07K 7/10, C12P 21/00,  
C12N 15/00, A61K 37/02,  
C07K 13/00

(30) Priority: 10.11.88 US 269583

(43) Date of publication of application:  
16.05.90 Bulletin 90/20

(94) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

(96) Date of deferred publication of the search report:  
30.01.91 Bulletin 91/05

(71) Applicant: MERCK & CO. INC.NC.  
126, East Lincoln Avenue P.O. Box 2000  
Rahway New Jersey 07065-0900(US)

Applicant: THE TRUSTEES OF THE  
UNIVERSITY OF PENNSYLVANIA  
c/o Office of Research Administration 133  
South 36th Street Suite 300  
Philadelphia Pennsylvania 19104-3246(US)

(72) Inventor: Shattil, Sanford J.  
510 Merwyn Road  
Narberth Pennsylvania 19072(US)  
Inventor: Taub, Rebecca A.  
403 Great Spring Road  
Rosemont Pennsylvania 19010(US)  
Inventor: Friedman, Paul A.  
403 Great Spring Road  
Rosemont Pennsylvania 19010(US)

(74) Representative: Stebbing, Peter John Hunter  
Stebbing & Co. 45 Lancaster Mewsews  
Lancaster Gate, London W2 3QQ(GB)

(54) Novel inhibitors of platelet binding.

(57) Peptides comprising the tripeptide sequence of arginine-tyrosine-aspartic acid, linked from amino- to carboxy-terminus, are inhibitors of fibrinogen binding, platelet aggregation, glycoprotein IIb-IIIa binding and related aberrant and normal physiological conditions.

**EP 0 368 486 A3**



European Patent  
Office

**PARTIAL EUROPEAN SEARCH REPORT**  
which under Rule 45 of the European Patent Convention  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

Application number  
**EP 89 31 0620**

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	THE JOURNAL OF CELL BIOLOGY, vol. 105, no. 5, November 1987, pages 2403-2408, The Rockefeller University Press; P.F. GOETINCK et al.: "The tandemly repeated sequences of cartilage link protein contain the sites for interaction with hyaluronic acid"  * Page 2404, column 1 *	1,3,4,5,6	C 07 K 5/08 C 07 K 7/06 C 07 K 7/10 C 12 P 21/00 C 12 N 15/00 A 61 K 37/02 C 07 K 13/00
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 264, no. 1, 5th January 1989, pages 259-265, The American Society for Biochemistry and Molecular Biology, Inc.; R. TAUB et al.: "A monoclonal antibody against the platelet fibrinogen receptor contains a ./. .		
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 K C 12 P C 12 N A 61 K
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-11,14-20 Claims searched incompletely: 12,13 Claims not searched: 12,13 Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention)</p>			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>07-11-1990</b>	Examiner <b>NOVOA Y SANJURJO</b>
<b>CATEGORY OF CITED DOCUMENTS</b>		<b>T</b> : theory or principle underlying the invention <b>E</b> : earlier patent document, but published on, or after the filing date <b>D</b> : document cited in the application <b>L</b> : document cited for other reasons  <b>A</b> : member of the same patent family, corresponding document	
<b>X</b> : particularly relevant if taken alone <b>Y</b> : particularly relevant if combined with another document of the same category <b>A</b> : technological background <b>O</b> : non-written disclosure <b>P</b> : intermediate document			



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application number

EP 89 31 0620

-2-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	sequence that mimics a receptor recognition domain in fibrinogen"  * Whole article *  --	1-20	
D,A	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, no. 26, 15th September 1988, pages 12948-12953, The American Society for Biochemistry and Molecular Biology, Inc.; J.S. BENNETT et al.: "Interaction of fibrinogen with its platelet receptor"  --		TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
D,A	THE JOURNAL OF BIOLOGICAL CHEMISTRY, 15th September 1985, pages 11107-11114, The American Society of Biological Chemists, Inc.; S.J. SHATTIL et al.: "Changes in the platelet membrane glycoprotein Iib-IIia complex during platelet activation"  ----		